

Symbols of cancer, do you recognized THEM? If yes, than join CFA in the fight, keep yourself healthy!

Let us talk about a very terrible disease that is commonly knew in our AFRICE community today, that rages from young kids to adults. Cancer doesn't see what a beautiful baby/child you are or dose it see what a wonder person you have bean in your lifetime! Cancer still take you from your love ones, this diseases have become one of the number one killer in the Africa community. Let us all say enough to cancer and STAND AGAINST it, the sooner you prevent yourself, is better for life. With insurance or not you can stay get tested by contracting caring for all 6000 bass lake rd crystal MN 55429 call 763- 537-0030 or 763-218-8685. To schedule a doctor appointment for you. cancer prevention is very important. If you have a specific cancel in mine that you would like to no more about please call us or write to us, if you like to remain anonymous too, we encourage you to start open talk in your home and community today about CANCER.



The analysis also revealed that the *lung cancer death rate* among those who have never *Nonsmoking black men* and women were found to have a higher death *rate*.



African-American women with advanced invasive epithelial ovarian carcinoma were less often treated with combined surgery and chemotherapy and more often treated with chemotherapy only. African-American women were twice as likely as white women not to receive appropriate treatment.



Synovial sarcoma is rare. It accounts for between 5 and 10 percent of the approximately 10,000 new soft tissue sarcomas reported each year (1). Synovial sarcoma occurs mostly in young adults, with a median age of 26.5 (1). Approximately 30 percent of patients with synovial sarcoma are younger than 20. This disease occurs more often in men than in women. About 50 percent of synovial sarcomas develop in the legs, especially synovial cancer to recur (come the knees. The second most common location is the arms (2). Less frequently, the disease develops in the trunk, head and neck region, or the abdomen (1, 2). It is common for back), usually within the first two years after treatment. Half of the cases of synovial sarcoma metastasize (spread to other areas of the body) to the lungs, lymph nodes, or bone marrow.



Prostate cancer is a significant health problem for middle-aged and elderly men. In the United States (US), it is the most frequently diagnosed cancer and is the second leading cause of cancer death. While men of all racial and ethnic backgrounds are at risk, black men of African descent are at especially high risk. African-Caribbean men, particularly Jamaican men, have the highest rate of prostate cancer in the world. The term African-American has been used to describe all black people living in the US. Use of such broad categorization ignores the existence of subcultures within the black community. While members of the black race may share similar primary, genetic characteristics, skin color cannot be equated with attitudes, knowledge, and behaviors of particular cultural groups. Therefore, pros ate cancer interventions developed for African-American men may not be effective for men of African-Caribbean descent.



**Leukemia and Lymphoma cancer, I am the killer that take the life's of innocent children and leave pain left in the heart of mothers. With the question why mine innocent child?**



2.7% in all women., 6.7% in black women., 7 deaths per 100,000 population per year (white). 12 deaths per 100,000 population per year (black). 4,600 deaths expected was in 2000 now is even more Mortality rates are about 50-80% lower than incidence rates. Black women have the highest mortality rate.



The *testicular cancer rate* has more than doubled among white *men* in the past 40 years, but has only recently begun to increase among *black men*.



**Breast Cancer If you are a lady please prevent me with regular checkup and joining the fight.**



Colon cancer, black men are diagnosed at more advanced stages of colon cancer than whites men, the report's authors note, and blacks have lower survival rates at each stage of diagnosis of most types of cancers



**Lymphoma cancel is a general term for several types of Hodgkin . Obesity and cancer risk among white and black United States.**



**Pancreatic cancer** The incidence *rate of pancreatic cancer* in the US *black population* is 10.8 cases per 100000 in whites. *Black men* also have the highest mortality *rate*



Men's breast cancer is something that few people are aware of, probably due to it accounting for only around 1% of all breast cancers. In 2007 it was estimated that in America men's breast cancer cases totaled approximately 2000 compared to about 178,000 for women. Out of those 2000 cases of men's breast cancer the survival rate is comparable with that of women suffering from breast cancer and depends largely on when it was diagnosed. Men of any age can develop breast cancer, but it is usually detected in men aged between 60 and 70.



**Melanoma/Skin Cancer** While there is no question that people of color are less likely to become afflicted with skin cancer, they are much more likely to die from it due to a delay in detection," said Perry Robins, M.D., President, The Skin Cancer Foundation. "Therefore, we feel it's our job to make these populations aware of the importance of early detection, prompt treatment and effective prevention."



Brain cancer can occur at any age. Studies have shown that two major age groups are affected. From ages 3 to 12 and 40 to 70 are the age groups when brain cancer is formed. Since researchers have been able to gather this data, it has led to the discovery of some risk factors. Workers in certain industries are at a higher risk for brain cancer than workers in other industries. These include, rubber manufacturing, drug manufacturing, and oil refining. Since brain cancer often occurs with members of the same family, heredity is believed to be another cause of brain cancer.



**The late Mr. Othello Garley**

The information here is provided for educational and informational purposes only, and is not intended to be a substitute for the diagnosis, treatment and advice of a qualified licensed medical professional. Please do not delay in seeking advice from a qualified licensed medical professional about treatment if you believe or feel you may have concerns about cancel